

Mathematical and Analytical Techniques with
Applications to Engineering

Oleg Kupervasser

Pole Solutions for Flame Front Propagation

 Springer

Attention!

This book can be sold here:

http://www.amazon.com/Propagation-Mathematical-Analytical-Applications-Engineering/dp/3319188445/ref=sr_1_1?ie=UTF8&qid=1430478768&sr=8-1&keywords=kupervasser

If you buy the book, please, write customer review. To do it, please, push yellow button “Write a customer review” in the end of the Internet page.

Mathematical and Analytical Techniques with Applications to Engineering

The importance of mathematics in the study of problems arising from the real world, and the increasing success with which it has been used to model situations ranging from the purely deterministic to the stochastic, in all areas of today's Physical Sciences and Engineering, is well established. The progress in applicable mathematics has been brought about by the extension and development of many important analytical approaches and techniques, in areas both old and new, frequently aided by the use of computers without which the solution of realistic problems in modern Physical Sciences and Engineering would otherwise have been impossible. The purpose of the series is to make available authoritative, up to date, and self-contained accounts of some of the most important and useful of these analytical approaches and techniques. Each volume in the series will provide a detailed introduction to a specific subject area of current importance, and then will go beyond this by reviewing recent contributions, thereby serving as a valuable reference source.

More information about this series at <http://www.springer.com/series/7311>

Oleg Kupervasser

Pole Solutions for Flame Front Propagation

 Springer

Oleg Kupervasser
Transist Video LLC
Moscow
Russia

ISSN 1559-7458 ISSN 1559-7466 (electronic)
Mathematical and Analytical Techniques with Applications to Engineering
ISBN 978-3-319-18844-7 ISBN 978-3-319-18845-4 (eBook)
DOI 10.1007/978-3-319-18845-4

Library of Congress Control Number: 2015941861

Springer Cham Heidelberg New York Dordrecht London
© Springer International Publishing Switzerland 2015

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made.

Printed on acid-free paper

Springer International Publishing AG Switzerland is part of Springer Science+Business Media
(www.springer.com)

Preface

This book deals with solving mathematically the unsteady flame propagation equations. New original mathematical methods for solving complex nonlinear equations and investigating their properties are presented. Pole solutions for flame front propagation are developed. Premixed flames and filtration combustion have remarkable properties: the complex nonlinear integro-differential equations for these problems have exact analytical solutions described by the motion of poles in a complex plane. Instead of complex equations, a finite set of ordinary differential equations is applied. These solutions help to investigate analytically and numerically properties of the flame front propagation equations.

The author solves mathematically unsteady flame propagation equations, describes new original methods for solving complex nonlinear equations and investigating their properties, and addresses open problems existing in the field of flame front propagation.

The theories (Laplacian growth, filtration combustion, premixed flame propagation) described in the book have many engineering and scientific applications, for example: investigation of burning in Ia supernovae; designing root modules and control of transport processes in space greenhouses; NASA Announces Microgravity Combustion Science Research Grants in Filtration Combustion for Microgravity Applications in 2002; Microgravity Fluids and Combustion Research was carried out at NASA Glenn Research Center; Microgravity smolder spread over a thin cellulosic fuel was studied with Radiative Ignition and Transition to Spread Investigation (RITSI) apparatus in the Glovebox Facility on the STS-75 USMP-3 space shuttle mission.

In Chap. 1 we describe basic concepts and problems in flame front propagation; give a review of many relevant papers and books in this field.

In Chap. 2 we investigate the problem of flame propagation. This problem is studied as an example of unstable fronts that wrinkle on many scales. The analytic tool of pole expansion in the complex plane is employed to address the interaction of the unstable growth process with random initial conditions and perturbations. We argue that the effect of random noise is immense and that it can never be neglected in sufficiently large systems. We present simulations that lead to scaling laws for the

velocity and acceleration of the front as a function of the system size and the level of noise, and analytic arguments that explain these results in terms of the noisy pole dynamics. We give a detailed description of excess number of poles in system, number of poles that appear in the system in unit of time, lifetime of pole. It allows us to understand dependence of the system parameters on noise.

In Chap. 3 we consider flame front propagation in channel geometries. The steady-state solution in this problem is space dependent, and therefore the linear stability analysis is described by a partial integro-differential equation with a space-dependent coefficient. Accordingly, it involves complicated eigenfunctions. We show that the analysis can be performed to required detail using a finite order dynamical system in terms of the dynamics of singularities in the complex plane, yielding detailed understanding of the physics of the eigenfunctions and eigenvalues.

In Chap. 4 flame propagation is used as a prototypical example of expanding fronts that wrinkle without limit in radial geometries, but reach a simple shape in channel geometry. We show that the relevant scaling laws that govern the radial growth can be inferred once the simpler channel geometry is understood in detail. In radial geometries (in contrast to channel geometries) the effect of external noise is crucial in accelerating and wrinkling the fronts. Nevertheless, once the interrelations between system size, velocity of propagation and noise level are understood in channel geometry, the scaling laws for radial growth follow.

In Chap. 5 Filtration Combustion is described by Laplacian growth without surface tension. These equations have elegant analytical solutions that replace the complex integro-differential motion equations by simple differential equations of pole motion in a complex plane. The main problem with such a solution is the existence of finite time singularities. To prevent such singularities, nonzero surface tension is usually used. However, nonzero surface tension does not exist in filtration combustion, and this destroys the analytical solutions. However, a more elegant approach exists for solving the problem. First, we can introduce a small amount of pole noise to the system. Second, for regularization of the problem, we throw out all new poles that can produce a finite time singularity. It can be strictly proved that the asymptotic solution for such a system is a single finger. Moreover, the qualitative consideration demonstrates that a finger with $1/2$ of the channel width is statistically stable. Therefore, all properties of such a solution are exactly the same as those of the solution with nonzero surface tension under numerical noise. The solution of the Saffman-Taylor problem without surface tension is similar to the solution for the equation of cellular flames in the case of the combustion of gas mixtures.

In Chap. 6 we give a short summary of the book ideas.

Moscow

Oleg Kupervasser

Acknowledgments

I would like to thank Zeev Olami, Barak Glanti, Itamar Procaccia, Mark Mineev-Weinstein, Leonid Kagan, Nikolay Nikolaevich Smirnov for their fruitful job and ideas, which were very useful for creating the book. I thank Editor Claus Ascheron for his very useful remarks and reviewing. I also thank Alexei Stanislavovich Shamaev for organization of very useful seminars and discussions in Moscow State University.

Contents

1	General View of Pole Solutions in Flame Front Propagation	1
2	Pole-Dynamics in Unstable Front Propagation: The Case of The Channel Geometry	11
2.1	Introduction.	11
2.2	Equations of Motion and Pole-Decomposition in the Channel Geometry	13
2.3	Linear Stability Analysis in Channel Geometry	15
2.3.1	Fourier Decomposition and Eigenvalues	16
2.3.2	Qualitative Understanding Using Pole-Analysis	18
2.3.3	Dynamics Near Marginality	18
2.3.4	Excitable System.	20
2.4	Initial Conditions, Pole Decomposition and Coarsening	21
2.4.1	Pole Expansion: General Comments.	21
2.4.2	The Initial Stages of the Front Evolution: The Exponential Stage and the Inverse Cascade.	23
2.4.3	Inverse Cascade in the Presence of Noise	26
2.5	Acceleration of the Premixed Flame Front, Pole Dynamics and Noise	27
2.5.1	Noisy Simulations	28
2.5.2	Calculation of the Number of Poles in the System	31
2.5.3	Theoretical Discussion of the Effect of Noise	35
2.5.4	The Acceleration of the Premixed Flame Front Because of Noise.	43
2.6	Comparison with Experiment for Premixed Flame in Channel	45
2.7	Summary and Conclusions	46
	Appendix: Derivation of Michelson Sivashinsky Equation.	47

3	Using of Pole Dynamics for Stability Analysis of Premixed Flame Fronts: Dynamical Systems Approach in the Complex Plane	55
3.1	Introduction.	55
3.2	Linear Stability Analysis in Channel Geometry	56
3.3	Linear Stability in Terms of Complex Singularities	58
3.3.1	The Modes Associated with the Giant Cusp	59
3.3.2	Modes Related to Additional Poles	61
3.4	Conclusions	65
4	Dynamics and Wrinkling of Radially Propagating Fronts Inferred from Scaling Laws in Channel Geometries	67
4.1	Introduction.	67
4.2	The Geometry of Developing Premixed Flame Fronts: Analysis with Pole Decomposition	68
4.3	Comparison with Experiment for Expanded Premixed Flames	77
4.4	Conclusions	83
5	Laplacian Growth Without Surface Tension in Filtration Combustion: Analytical Pole Solution	85
5.1	Introduction.	85
5.2	Asymptotic Single Saffman-Taylor “finger” Formation Without Surface Tension.	88
5.2.1	Asymptotic Behaviour of the Poles in the Mathematical Plane.	90
5.2.2	Theorem About Coalescence of the Poles	94
5.2.3	The Final Result	95
5.3	Saffman-Taylor “finger” Formation with Half of the Channel Size	97
5.4	Comparison Laplacian Growth with Experiment for Filtration Combustion	102
5.5	Conclusions	104
	Appendix 1: Proof of Theorems.	104
	Appendix 2: Filtration Combustion—Basic Equations.	105
6	Summary	109
	Bibliography	111
	Index	117

Chapter 1

General View of Pole Solutions in Flame Front Propagation

Abstract We describe basic concepts and problems in flame front propagation; give a review of many relevant papers and books in this field.

Problems of interface growth have received much attention [1–11]. Such are, for example, the diffusion limited aggregation (DLA) [12], random sequential adsorption (RSA) [13], Laplacian growth [6, 14–23] or premixed flame front propagation [1, 3, 4, 7, 24–29]. We will mainly pay attention in this book to the numerical and analytical investigation of the last two problems.

We start to study the example of premixed flame propagation rather than Laplacian growth, simply because the former has an analytic description in terms of poles also in the experimentally relevant case of finite regularization term.

In addition to the fact that premixed flame front propagation is an interesting physical problem we feel that we can also explain experimental results on the basis of theoretical investigations.

The premixed flame—the self-sustaining wave of an exothermic chemical reaction—is one of the basic manifestations of gaseous combustion. It is well established, however, that the simplest imaginable flame configuration—unbounded planar flame freely propagating through initially motionless homogeneous combustible mixture—is intrinsically unstable and spontaneously assumes a characteristic two- or three-dimensional structure.

This book considers the very interesting problem of describing the nonlinear stage of development of hydrodynamic instability of the premixed flame. This problem can be considered for 1D (channel propagation), 2D (cylindrical propagation or rectangular channel cases), 3D (spherical case). The dimensionality of the problem is defined by the space dimensionality of the range of definition for the function, describing moving front. The direct numerical simulations can be made based on of the Navier-Stokes equations for expanding premixed flames [30–33] and for channel geometry [34–36]. For 2D-cylindrical and 3D cases we can see experimentally observable effects [37–45] of the self-acceleration of the front of a divergent premixed flame, the formation of cellular structure, and other effects.

Much more simple equation for premixed flame can be obtained [29, 46]. It is Michelson-Sivashinsky approximation model for the channel case (and very similar

Sivashinsky, Filyand and Frankel equation for cylindrical case). The Michelson-Sivashinsky model assumes very serious limitations, such as the smallness of the coefficient of gas expansion and, consequently, the potential flow in the combustion products and fresh mixture, weak nonlinearity, the assumptions of the stabilizing effect of the curvature of the premixed flame front and a linear dependence on the curvature of the normal speed, and others.

Interest of this model stems, firstly, from the fact that despite the serious limitations, this model can qualitatively describe the scenario of hydrodynamic instability and, in particular, the self-acceleration of the front of a divergent premixed flame, the formation of a cellular structure, and other effects.

Secondary, this nonlinear model has exact solutions which can be constructed on the basis of pole expansions. This pole expansion method got development in the following papers [47–50]. There was investigated relationships between pole solutions and partial decomposition in Fourier series [51], between the gas flow field and pole solutions [52]. The future development can be found in the following papers and the correspondent references inside of [24–27]. The Michelson-Sivashinsky model and pole solutions can be used for investigation of burning in Ia supernovae [53].

The aim of this book is to examine the role of random fluctuations on the dynamics of growing wrinkled interfaces which are governed by non-linear equations of motion. We are interested in those examples for which the growth of a flat or smooth interface are inherently unstable. A famous example of such growth phenomena is provided by Laplacian growth patterns [6, 8–10, 14, 15, 18–20]. The experimental realization of such patterns is seen for example in Hele-Shaw cells [6, 10, 16] in which air or another low viscosity fluid is displacing oil or some other high viscosity fluid. Under normal conditions the advancing fronts do not remain flat; in channel geometries they form in time a stable finger whose width is determined by delicate effects that arise from the existence of surface tension. In radial geometry, the growth the interface forms a contorted and ramified fractal shape. The same model of Laplacian growth can be used for description of Filtration Combustion, where no surface tension exists [14, 15, 18–20] (see also Appendix 2 in Chap. 5). A related phenomenon has been studied in a model equation for premixed flame propagation [29] which has the same linear stability properties as the Laplacian growth problem. The physical problem in this case is that of premixed flames which exist as self-sustaining fronts of exothermic chemical reactions in gaseous combustion. Experiments on Premixed flame propagation in radial (cylindrical) [37] and spherical [37–45] geometry show that the premixed flame front accelerates as time goes on, and roughens with characteristic exponents. Both observations must receive proper theoretical explanations. It is notable that the channel and radial growth are markedly different; the former leads to a single giant cusp in the moving front, whereas the latter exhibits infinitely many cusps that appear in a complex hierarchy as the premixed flame front develops ([28, 46] and Chap. 4).

Analytic techniques to study such processes are available [47]. In the context of premixed flame propagation [24–28, 48–50], and in Laplacian growth in the zero surface tension limit [14, 16, 22, 54, 55] one can examine solutions that are described in terms of poles in the complex plane. This description is very useful in providing a

set of ordinary differential equations for the positions of the poles, from which one can deduce the geometry of the developing front in an extremely economical and efficient way. Unfortunately this description is not available in the case of Laplacian growth with surface tension, and this makes the premixed flame propagation problem very attractive. However, it suffers from one fundamental drawback. For the noiseless equation the pole-dynamics always conserves the number of poles that existed in the initial conditions. As a result there is a final degree of ramification that is afforded by every set of initial conditions even in the radial geometry, and it is not obvious how to describe the continuing self-similar growth that is seen in experimental conditions or numerical simulations. Furthermore, as mentioned before, at least in the case of premixed flame propagation one observes [37–45] an *acceleration* of the premixed flame front with time. Such a phenomenon is impossible when the number of poles is conserved. It is therefore tempting to conjecture that noise may have an important role in affecting the actual growth phenomena that are observed in such systems. In fact, the effect of noise on unstable front dynamics must be adequately described. From the point of view of analytic techniques noise can certainly generate new poles even if the initial conditions had a finite number of poles. The subject of pole dynamics with the existence of random noise, and the interaction between random fluctuations and deterministic front propagation are the main issues of this book.

The consideration of the noise term is critically important for these processes. Indeed, in papers [56–60], the authors try to explain self-acceleration without involvement of the external forcing. However, no self-acceleration exist for the finite number of poles. So we can explain the self-acceleration and the appearance of new poles or by the noise or by the “rain” of poles from the “cloud” in infinity. Unfortunately, such rain can not “achieve” the “ground” for accelerating premixed flame as we will see in Chap. 4.

The authors of [56–60] suppose that some analytical solution without noise corresponds to the unsteady solution with increasing number of cusps in radial case. They base this conclusion on the weak dependence of numerical simulations on the noise reduction ([56], Fig. 2). However, we will see in Chap. 2 that for the big interval of the noise values (larger than some small value and up to some big value) in the regime III (corresponding to the increasing number of poles in radial case), the dependence on the noise is very slow: $f^{0.02}$. This result explains the weak dependence of numerical simulations on the noise reduction.

The calculations with a noise term was performed in both one-dimensional and two-dimensional formulations of the problem in the papers [24–27] and by Karlin and Sivashinsky [61, 62] for 1D, 2D and 3D cases.

The future development of these results is gotten in papers [63–65]. In the paper [63] the noise term was considered in the poles-like form. The numerical results for 1D case are in a good consistence with the theoretical results of this book. The acceleration value of premixed flames fronts changes in the interval 1.25–1.5 according experimental [37–45] and theoretical [24, 26, 27, 30, 33, 46, 61–63, 80–84] results. The dependence of the acceleration value on the other parameters can explain this variation. The dependence was investigated numerically [63] and experimentally

[40]. In [65], hydrodynamic instability of inward-propagating premixed flames was investigated.

Joulin et al. [66–69] use a vary similar to [24–28] approach for the channel and radial premixed flame growth. Main attention in the channel case in Joulin’s work was made to the investigation of mean-spacing between cusps (crests). For the radial case only the linear dependence of radius on time (no self-acceleration) is considered in Joulin’s work. The main attention in our work is made to the velocity of premixed flame front (self-acceleration for the radial case) and the premixed flame front width.

In the next papers of Joulin et al. [50, 70, 71], the pole solutions for the Zhdanov-Trubnikov equation (extensions of Michelson-Sivashinsky equation that incorporate higher orders of the coefficient of gas expansion) are developed.

In the paper of Joulin et al. [72] the pole solutions for steady forced premixed flames are considered.

In the paper of Joulin et al. [73], the steady pole solutions is analyzed for Neumann boundary conditions (“no-flux” boundary conditions suggested in [74, 75], which are more physical than periodical boundary conditions [48]) in the Michelson-Sivashinsky equation. For these boundary conditions, we get two cusps on boundaries with unequal fluctuating sizes (instead of one moving giant cusp for periodical boundary conditions).

Similar “no-flux” boundary conditions (instead periodical ones [22, 23]) was suggested previously for Laplacian growth in [76] and developed in [14, 21]. However, [76] repeats the conclusion about one finger asymptotic, which are made previously in [16].

In the paper of Joulin et al. [49, 77] approximate the giant cusp steady solution of the Michelson-Sivashinsky equation by parabola (geometrical stretch) and investigate its nonlinear stability: perturbations of this approximated solution can be described by poles.

In the paper of Joulin et al. [78], the authors discovered that pole positions in the giant cusp (for the number of poles $N > 3$) can be approximated with high precision by roots of polynomials, ruled by a discretized Burgers equation.

In the paper of Gostintsev, Istratov and Shulenin [37] an interesting survey of experimental studies on outward propagating spherical and cylindrical premixed flames in the regime of well developed Darrieus-Landau hydrodynamic instability [79] is presented. The available data clearly indicate that freely expanding wrinkled premixed flames possess two intrinsic features:

1. Multi-quasi-cusps structure of the premixed flame front. (The premixed flame front consists of a large number of quasi-cusps, i.e., cusps with rounded tips.)
2. Noticeable acceleration of the premixed flame front: the propagation of wrinkled premixed flames really self accelerates and the acceleration follows a self-similar law. In other words, we fit the velocity data with a power-law formula, the exponent not only is greater than unity but it is also a constant, at least within a certain stage of the propagation.

According [37], the temporal dependence of the premixed flame radius is nearly identical for all premixtures discussed and correlates well with the simple relation:

$$R_0(t) = At^{3/2} + B \quad (1.1)$$

Here $R_0(t)$ is the effective (average) radius of the wrinkled premixed flame and A, B are empirical constants.

In follow-on publications (both this authors and the other authors [37–45]) the value of acceleration 1.5 was lowered to 1.25–1.5, with the revised assessment that the value of 1.5 is only attained in limiting cases.

In this book we study the spatial and temporal behavior of a nonlinear continuum model (i.e., a model which possesses an infinite number of degrees of freedom) which embodies all the characteristics deemed essential to premixed flame systems; namely, dispersiveness, nonlinearity and linear instability. Sivashinsky, Filyand and Frankel [46] obtained an equation, denoted by SFF in what follows, to describe how two-dimensional wrinkles of the cylindrical premixed flame grow as a consequence of the well-known Darrieus-Landau hydrodynamic instability [79]. The formal derivation of (1.2) is quite similar to that Michelson-Sivashinsky (MS) (1.5) presented by Sivashmsky [29], where the hydrodynamically unstable planar premixed flames were studied. The SFF and MS equations were obtained under assumption that the Atwood number $A = (E - 1)/(E + 1)$ (based on the fresh-to-burnt density ratio $E > 1$) is small.

The SFF equation reads as follows:

$$\frac{\partial R}{\partial t} = \frac{U_b}{2R_0^2(t)} \left(\frac{\partial R}{\partial \theta} \right)^2 + \frac{D_M}{R_0^2(t)} \frac{\partial^2 R}{\partial \theta^2} + \frac{\gamma U_b}{2R_0(t)} I\{R\} + U_b. \quad (1.2)$$

where $0 < \theta < 2\pi$ is an angle, $R(\theta, t)$ is the modulus of the radius-vector on the premixed flame interface (We can to increase the define area of the function $R(\theta, t)$ with periodical boundary condition to $-\infty < \theta < +\infty$ by the periodical continuation), U_b, D_M, γ are constants.

$$\begin{aligned} I(R) &= \frac{1}{\pi} \sum_{n=1}^{\infty} n \int_0^{2\pi} \cos[n(\theta - \theta^*)] R(\theta^*, t) d\theta^* = \\ &= -\frac{1}{\pi} P \int_{-\infty}^{+\infty} \frac{\frac{\partial R(\theta^*, t)}{\partial \theta^*}}{\theta^* - \theta} d\theta^* \end{aligned} \quad (1.3)$$

$$R_0(t) = \frac{1}{2\pi} \int_0^{2\pi} R(\theta, t) d\theta. \quad (1.4)$$

Sivashinsky, Filyand and Frankel [46] made a direct numerical simulation of this nonlinear evolution equation for the cylindrical premixed flame interface dynamics.

The result obtained shows that the two mentioned experimental effects take place. Moreover, the evaluated acceleration rate is not incompatible with the power law given by (1.1). For comparison, numerical simulations of freely expanding diffusively

unstable premixed flames were presented as well. In this case no tendency towards acceleration has been observed.

The acceleration value of premixed flames fronts changes in the interval 1.25–1.5 according the other theoretical results [24, 26, 27, 30, 33, 61–63, 80–84].

To obtain results for radial premixed flame growth it is necessary [26, 30, 85] to investigate the channel case first. The channel version of equation for premixed flame front propagation is the so-called Michelson-Sivashinsky equation [29, 48, 86] (see also Appendix in Chap. 2) and looks like

$$\frac{\partial H}{\partial t} = \frac{1}{2} \left(\frac{\partial H}{\partial x} \right)^2 + \nu \frac{\partial^2 H}{\partial x^2} + I\{H\}. \quad (1.5)$$

$$I(H) = -\frac{1}{\pi} P \int_{-\infty}^{+\infty} \frac{\frac{\partial H(x^*, t)}{\partial x^*}}{x^* - x} dx^*. \quad (1.6)$$

with periodic boundary condition on the interval $x \in [0, L]$, where L is size of the system. ν is constant, $\nu > 0$. H is the height of the premixed flame front point, $P \int$ is the usual principal value integral.

There exists possibility to use methods found for the premixed flame front propagation, in different fields where similar problems appear such as the important model of Laplacian growth.

In the absence of surface tension, whose effect is to stabilize the short-wavelength perturbations of the interface, the problem of 2D Laplacian growth is described as follows

$$(\partial_x^2 + \partial_y^2)u = 0. \quad (1.7)$$


$$u|_{\Gamma(t)} = 0, \quad \partial_n u|_{\Sigma} = 1. \quad (1.8)$$

$$v_n = \partial_n u|_{\Gamma(t)}. \quad (1.9)$$

Here $u(x, y; t)$ is the scalar field mentioned, $\Gamma(t)$ is the moving interface, Σ is a fixed external boundary, ∂_n is a normal component of the gradient to the boundary (i.e. the normal derivative), and v_n is a normal component of the velocity of the front.

Equations for premixed flame front propagation and Laplacian growth with zero surface tension have remarkable property : these equations can be solved in terms of poles in the complex plane [22, 47–49, 69, 87]. So we obtain a set of ordinary differential equations for the coordinates of these poles. The number of the poles is constant value in the system, but to explain such effect as growth of the velocity premixed flame front we need to consider some noise that is a source of new poles. So we need to solve the problem of interaction of the random fluctuations and the pole motion.

The simplest case is the channel geometry. Main results for this case is existence of the giant cusp solution [48] (Fig. 1.1), which is represented in configuration space

Fig. 1.1 Giant cusp solution

by poles which are organized on a line parallel to the imaginary axis. This pole solution is an attractor for pole dynamics.

A complete analysis of this steady-state solution was first presented in [48] and the main results are summarized as follows:

1. There is only one stable stationary solution which is geometrically represented by a giant cusp (or equivalently one finger) and analytically by $N(L)$ poles which are aligned on one line parallel to the imaginary axis. The existence of this solution is made clearer with the following remarks.
2. There exists an attraction between the poles along the real line. The resulting dynamics merges all the x positions of poles whose y -position remains finite.
3. The y positions are distinct, and the poles are aligned above each other in positions $y_{j-1} < y_j < y_{j+1}$ with the maximal being $y_{N(L)}$. This can be understood from equations for the poles motion in which the interaction is seen to be repulsive at short ranges, but changes sign at longer ranges.
4. If one adds an additional pole to such a solution, this pole (or another) will be pushed to infinity along the imaginary axis. If the system has less than $N(L)$ poles it is unstable to the addition of poles, and any noise will drive the system towards this unique state. The number $N(L)$ is

$$N(L) = \left[\frac{1}{2} \left(\frac{L}{\nu} + 1 \right) \right], \quad (1.10)$$

where $[\dots]$ is the integer part and $2\pi L$ is a system size. To see this consider a system with N poles and such that all the values of y_j satisfy the condition $0 < y_j < y_{max}$. Add now one additional pole whose coordinates are $z_a \equiv (x_a, y_a)$ with $y_a \gg y_{max}$. From the equation of motion for y_a , we see that the terms in the

sum are all of the order of unity as is also the $\cot(y_a)$ term. Thus the equation of motion of y_a is approximately

$$\frac{dy_a}{dt} \approx \nu \frac{2N + 1}{L^2} - \frac{1}{L}. \quad (1.11)$$

The fate of this pole depends on the number of other poles. If N is too large the pole will run to infinity, whereas if N is small the pole will be attracted towards the real axis. The condition for moving away to infinity is that $N > N(L)$ where $N(L)$ is given by (1.10). On the other hand the y coordinate of the poles cannot hit zero. Zero is a repulsive line, and poles are pushed away from zero with infinite velocity. To see this consider a pole whose y_j approaches zero. For any finite L the term $\coth(y_j)$ grows unboundedly whereas all the other terms in the equation for the poles motion remain bounded.

5. The height of the cusp is proportional to L . The distribution of positions of the poles along the line of constant x was worked out in [48].

We will refer to the solution with all these properties as the Thual-Frisch-Henon TFH-cusp solution.

The main results of our own work are as follow. Traditional linear analysis was made for this giant cusp solution. This analysis demonstrates the existence of negative eigenvalues that go to zero when the system size goes to infinity.

1. There exists an obvious Goldstone or translational mode with eigenvalue $\lambda_0 = 0$. This eigenmode stems from the Galilean invariance of the equation of motion.
2. The rescaled eigenvalues ($L^2 \lambda_i$) oscillate periodically between values that are L -independent in this presentation. In other words, up to the oscillatory behavior the eigenvalues depend on L like L^{-2} .
3. The eigenvalues λ_1 and λ_2 hit zero periodically. The functional dependence in this presentation appears almost piecewise linear.
4. The higher eigenvalues also exhibit similar qualitative behavior, but without reaching zero. We note that the solution becomes marginally stable for every value of L for which the eigenvalues λ_1 and λ_2 hit zero. The L^{-2} dependence of the spectrum indicates that the solution becomes more and more sensitive to noise as L increases.

It was proved that arbitrary initial conditions can be written in the term of poles in the complex plane. Inverse cascade process of giant cusp formation was investigated numerically and analytically. Dependencies of the premixed flame front width and mean velocity were found. The next step in investigation of the channel case was the influence of random noise on the pole dynamics. The main effect of the external noise is the appearance of new poles in the minima of the premixed flame front and the merging these poles with the giant cusp. The dependence of the mean premixed flame front velocity on the noise and the system size was found. The velocity is almost independent on the noise until the noise achieves some critical value. In the

dependence of the velocity on the system size we see growth of the velocity with some exponent until the velocity achieves some saturation value.

Denoting v as the velocity of the premixed flame front and L the system size:

1. We can see two different regimes of behavior the average velocity v as a function of noise f for fixed system size L . For the noise f smaller than some fixed value f_{cr}

$$v \sim f^\xi. \quad (1.12)$$

For these values of f this dependence is very weak, and $\xi \approx 0.02$. For large values of f the dependence is much stronger

2. We can see growth of the average velocity v as a function of the system size L . After some values of L we can see saturation of the velocity. For regime $f < f_{cr}$ the growth of the velocity can be written as

$$v \sim L^\mu, \quad \mu \approx 0.35 \pm 0.03. \quad (1.13)$$

3. For $f > f_{cr}$ in Fig. 3.9 we will see qualitative change in the appearance of the premixed flame front: the noise introduces significant levels of small scales structure in addition to the cusps. The same observation was made in [88]

The dependence of the number of poles in the system and the number of the poles that appear in the system in unit time was investigated numerically as a function of the noise and the system parameters. The life time of a pole was found numerically. Theoretical discussion of the effect of noise on the pole dynamics and mean velocity was made [24, 27].

Pole dynamics can be used also to analyze small perturbation of the premixed flame front and make the full stability analysis of the giant cusp. Two kinds of modes were found. The first one is eigenoscillations of the poles in the giant cusp. The second one is modes connected to the appearance of the new poles in the system. The eigenvalues of these modes were found in [25, 89, 90]. The results of these papers are in good agreement with the traditional stability analysis ([89, 90] repeat many conclusions made previously in [25]).

The results found for the channel case can be used to analyze premixed flame front propagation in the radial case [26, 28]. Main feature of this case is a competition between attraction of the poles and expanding of the premixed flame front. So in this case we obtain not only one giant cusp but a set of cusps. New poles that appear in the system because of the noise form these cusps. On the basis of the equation of poles motion we can find connection between acceleration of the premixed flame front and the width of the interface. On the basis of the result for mean velocity in the channel case the acceleration of the premixed flame front can be found. So we obtain full picture of the premixed flame front propagation in the radial case.

The next step in the investigation of the problem is considering Laplacian growth with zero surface tension that also has pole solutions. In the case of Laplacian growth we obtain result that is analogous to the merging of the poles in the channel case of

the premixed flame front propagation: all poles coalesce into one pole in the case of periodic boundary condition or two poles on the boundaries in the case of no-flux boundary conditions. This result can be proved theoretically [14, 16]. Moreover, it is possible to demonstrate that this solution is statistically stable: the width of the final finger will be oscillate near $1/2$ of the channel width in the presence of a noise [14].

The structure of this book is as follow.

In Chap. 2 we obtain main results for the channel case of the premixed flame front propagation. We give results about steady state solutions, present traditional linear analysis of the problem and investigate analytically and numerically the influence of noise on the mean velocity of the front and pole dynamics.

In Chap. 3 we obtain results of the linear stability analysis by the help of pole solutions.

In Chap. 4 we use the result obtained for the channel case for analysis of the premixed flame front propagation in the radial case

In Chap. 5 we investigate asymptotic behavior of the poles in the complex plane for the Laplacian growth with the zero surface tension in the case of periodic and no-flux boundary condition.

In Chap. 6 we give a short summary of the book ideas.

Chapter 2

Pole-Dynamics in Unstable Front Propagation: The Case of The Channel Geometry

Abstract We investigate the problem of flame propagation. This problem is studied as an example of unstable fronts that wrinkle on many scales. The analytic tool of pole expansion in the complex plane is employed to address the interaction of the unstable growth process with random initial conditions and perturbations. We argue that the effect of random noise is immense and that it can never be neglected in sufficiently large systems. We present simulations that lead to scaling laws for the velocity and acceleration of the front as a function of the system size and the level of noise, and analytic arguments that explain these results in terms of the noisy pole dynamics. We make detailed description of excess number of poles in system, number of poles that appear in the system in unit of time, life time of pole. It allows us to understand dependence of the system parameters on noise.

2.1 Introduction

It must be mentioned that the simplest 1D case of the premixed flame front propagation is very important. It was the main reason for creation this chapter considering in detail the 1D case. Such investigation of this case allows us to understand qualitatively and quantitatively the pole dynamics. This understanding is a basis for the consideration more complex 2D and 3D cases. The papers [26, 30, 85] clearly demonstrate this fact. The cellular structure, acceleration exponent for 2D case was found on the basis of 1D results.

We choose to begin the study with channel geometry. The reason is that in radial geometry it is more difficult to disentangle the effects of external noise from those of initial conditions. After all, initially the system can contain infinitely many poles, very far away near infinity in the complex plane (and therefore having an infinitely small contribution to the interface). Since the growth of the radius changes the stability of the system, more and more of these poles might fall down to the real axis and become observable. In channel geometry the analysis of the effect of initial conditions is relatively straightforward, and one can understand it before focusing on the (more interesting) effects of external noise [48]. The basic reason for this is that in

Chapter 6

Summary

Abstract We give a short summary of the book ideas.

The problem of premixed flame propagation is studied as an example of unstable fronts that wrinkle on many scales. The analytic tool of pole expansion in the complex plane is employed to address the interaction of the unstable growth process with random initial conditions and perturbations. We argue that the effect of random noise is immense and that it can never be neglected in sufficiently large systems. We present simulations that lead to scaling laws for the velocity and acceleration of the front as a function of the system size and the level of noise, and analytic arguments that explain these results in terms of the noisy pole dynamics.

We consider premixed flame front propagation in channel geometries. The steady state solution in this problem is space dependent, and therefore the linear stability analysis is described by a partial integro-differential equation with a space dependent coefficient. Accordingly it involves complicated eigenfunctions. We show that the analysis can be performed to required detail using a finite order dynamical system in terms of the dynamics of singularities in the complex plane, yielding detailed understanding of the physics of the eigenfunctions and eigenvalues.

The roughening of expanding premixed flame fronts by the accretion of cusp-like singularities is a fascinating example of the interplay between instability, noise and nonlinear dynamics that is reminiscent of self-fractalization in Laplacian growth patterns. The nonlinear integro-differential equation that describes the dynamics of expanding premixed flame fronts is amenable to analytic investigations using pole decomposition. This powerful technique allows the development of a satisfactory understanding of the qualitative and some quantitative aspects of the complex geometry that develops in expanding premixed flame fronts.

Premixed flame propagation is used as a prototypical example of expanding fronts that wrinkle without limit in radial geometries but reach a simple shape in channel geometry. We show that the relevant scaling laws that govern the radial growth can be inferred once the simpler channel geometry is understood in detail. In radial geometries (in contrast to channel geometries) the effect of external noise is crucial in accelerating and wrinkling the fronts. Nevertheless, once the interrelations between system size, velocity of propagation and noise level are understood in channel geometry, the scaling laws for radial growth follow.

The mathematical problem of Laplacian growth without surface tension exhibits a family of exact (analytic) solutions in terms of logarithmic poles in the complex plane. We show that this family of solutions has a remarkable property: generic initial conditions in channel geometry which begin with arbitrarily many features exhibit an inverse cascade into a single finger. Moreover, it is possible to demonstrate that this solution is statistically stable: the width of the final finger will be oscillate near $1/2$ of the channel width in the presence of a noise.

Bibliography

1. A. Lipatnikov, *Fundamentals of Premixed Turbulent Combustion* (CRC Press, Boca Raton, 2013)
2. M.T. Torres, C.R. Huerta, *Hybrid Filtration Combustion in Book: Hydrogen Energy Challenges and Perspectives*, ed. by D. Minic (InTech, Croatia, 2012)
3. N. Swaminathan, K. Bray (eds.), *Turbulent Premixed Flames* (Cambridge University Press, Cambridge, 2011)
4. M. Liberman, *Introduction to Physics and Chemistry of Combustion: Explosion, Flame, Detonation* (Springer, Berlin, 2008)
5. Z. Chen, G. Huan, Y. Ma, *Computational Methods for Multiphase Flows in Porous Media* (SIAM, Philadelphia, 2006)
6. B. Gustafsson, A. Vasil'ev, *Conformal and Potential Analysis in Hele-Shaw Cells*. Advances in Mathematical Fluid Mechanics (Basel, Birkhauser, 2006)
7. V.V. Bychkov, M.A. Liberman, Dynamics and stability of premixed flames. *Phys. Rep.* **325**, 115–237 (2000)
8. A.-L. Barbási, H.E. Stanley, *Fractal Concepts in Surface Growth* (Cambridge University Press, New York, 1995)
9. T. Viscek, *Fractal Growth Phenomena* (World Scientific, Singapore, 1992)
10. P. Pelce, *Dynamics of Curved Fronts* (Academic Press, Boston, 1988)
11. G.H. Markstein (ed.), *Nonsteady Flame Propagation* (Pergamon Press, Oxford, 1964)
12. T.A. Witten Jr, L.M. Sander, *Phys. Rev. Lett.* **47**, 1400 (1981)
13. J.W. Evans, *Rev. Mod. Phys.* **65**, 1281 (1993)
14. O. Kupervasser, Laplacian Growth without Surface Tension in Filtration Combustion: Analytical Pole Solution, Complexity (Wiley Periodicals Inc, 2014), <http://onlinelibrary.wiley.com/doi/10.1002/cplx.21627/abstract>. Accessed 4 Nov 2014
15. A.P. Aldushin, B.Sh. Braverman, Saffman Taylor problem in filtration combustion. *Russ. J. Phys. Chem. B* **4**(5), 788–792 (2010)
16. M. Mineev-Weinstein, O. Kupervasser, *Formation of a Single Saffman-Taylor Finger after Fingers Competition: An Exact Result in the Absence of Surface Tension*. 82nd Statistical Mechanics Meeting, Rutgers University, 10–12 December 1999 (*J. Stat. Phys.* **99**(3–4), 1057–1065 (2000))
17. B.J. Matkowsky, Filtration combustion in smoldering and SHS, in *Fifth International Microgravity Combustion Workshop*, Cleveland, Ohio (1999)
18. A.P. Aldushin, B.J. Matkowsky, Extremum principles for selection in the Saffman-Taylor finger and Taylor-Saffman bubble problems. *Phys. Fluids* **11**, 1287–1296 (1999)

19. A.P. Aldushin, B.J. Matkowsky, Instabilities, fingering and the Saffman-Taylor problem in filtration combustion. *Combust. Sci. Technol.* **133**, 293–341 (1998)
20. A.P. Aldushin, B.J. Matkowsky, Selection in the Saffman-Taylor finger problem and the Taylor-Saffman bubble problem without surface tension. *Appl. Math. Lett.* **11**, 57–62 (1998)
21. M. Mineev-Weinstein, *Phys. Rev. Lett.* **80**, 2113–2116 (1998)
22. S. Ponce Dawson, M. Mineev-Weinstein, *Physica D* **73**, 373–387 (1994)
23. M. Mineev-Weinstein, S. Ponce Dawson, *Phys. Rev. E* **50**, R24 (1994)
24. O. Kupervasser, Z. Olami, Random noise and pole-dynamics in unstable front propagation. *Combust. Explos. Shock Waves* **49**(3), 141–152 (2013)
25. O. Kupervasser, Z. Olami, I. Procaccia, Stability analysis of flame fronts: dynamical systems approach in the complex plane. *Phys. Rev. E* **59**(3), 2587–2593 (1999)
26. B. Glanti, O. Kupervasser, Z. Olami, I. Procaccia, Dynamics and wrinkling of radially propagation fronts inferred from scaling law in channel geometries. *Phys. Rev. Lett.* **80**(11), 2477–2480 (1998)
27. Z. Olami, B. Glanti, O. Kupervasser, I. Procaccia, Random noise and pole-dynamics in unstable front propagation. *Phys. Rev. E* **55**(3), 2649–2663 (1997)
28. O. Kupervasser, Z. Olami, I. Procaccia, Geometry of developing flame fronts: analysis with pole decomposition. *Phys. Rev. Lett.* **76**, 146–149 (1996)
29. G.I. Sivashinsky, Nonlinear analysis of hydrodynamics instability in laminar flames I. Derivations fo basic equations. *Acta Astronautica* **4**, 1177–1206 (1977)
30. M.A. Liberman, M.F. Ivanov, O.E. Peil, D.M. Valiev, L.-E. Eriksson, Self-acceleration and fractal structure of outward freely propagating flames. *Phys. Fluids* **16**(7), 2476–2482 (2004)
31. E. Albin, Y. D'Ángelo, Assessment of the evolution equation modelling approach for three-dimensional expanding wrinkled premixed flames. *Combust. Flame Elsevier* **159**(5), 1932–1948 (2012)
32. F. Creta, M. Matalon, Strain rate effects on the nonlinear development of hydrodynamically unstable flames. *Proc. Combust. Inst.* **33**, 1087–1094 (2011)
33. Y.X. Xin, C.S. Yoo, J.H. Chen, C.K. Law, A DNS study of self-accelerating cylindrical hydrogenair flames with detailed chemistry. *Proc. Combust. Inst.* **35**(1), 753–760 (2015)
34. P.E. Hamlington, A.Y. Poludnenko, E.S. Oran, Interactions between turbulence and flames in premixed reacting flows. *Phys. Fluids* **23**, 125111 (2011)
35. C. Altantzis, C.E. Frouzakis, A.G. Tomboulides, S.G. Kerkemeier, K. Boulouchos, Detailed numerical simulations of intrinsically unstable two-dimensional planar lean premixed hydrogen/air flames. *Proc. Combust. Inst.* **33**, 1261–1268 (2011)
36. N. Fogla, F. Creta, M. Matalon, Influence of the Darrieus-Landau instability on the propagation of planar turbulent flames. *Proc. Combust. Inst.* **34**, 15091517 (2013)
37. Yu.A. Gostintsev, A.G. Istratov, Yu.V. Shulenin, Self-similar propagation of a free turbulent flame in mixed gas mixtures. *Combust. Explos. Shock Waves* **24**(5), 63–70 (1988)
38. F. Wu, G. Jomaas, C.K. Law, On self-acceleration of cellular spherical flames. Fall Technical Meeting of the Eastern States Section of the Combustion, Institute Hosted by the University of Connecticut, Storrs, CT (2011), <http://lcg.princeton.edu/publications/conference-publications/wu-on-self-acceleration-of-cellular-spherical-flames.aspx>
39. D. Bradley, T.M. Cresswell, J.S. Puttock, Flame acceleration due to flame-induced instabilities in large-scale explosions. *Combust. Flame* **124**, 551–559 (2001)
40. W.K. Kim, T. Mogi, K. Kuwanac, R. Dobashi, Self-similar propagation of expanding spherical flames in large scale gas explosions. *Proc. Combust. Inst.* **35**(2), 2051–2058 (2015)
41. F. Wu, G. Jomaas, C.K. Law, An experimental investigation on self-acceleration of cellular spherical flames. *Proc. Combust. Inst.* **34**, 937–945 (2013)
42. M.Z. Haq, Correlations for the onset of instabilities of spherical laminar premixed flames. *J. Heat Trans.* **127**, 1410 (2005)
43. O.C. Kwon, G. Rozenchan, C.K. Law, Cellular instabilities and self-acceleration of outwardly propagating spherical flames. *Proc. Combust. Inst.* **29**, 1775–1783 (2002)
44. Y.A. Gostintsev, A.G. Istratov, N.I. Kidin, V.E. Fortov, Self-turbulization of gas flames: an analysis of experimental results. *High Temp.* **37**(2), 282–288 (1999)

45. Y.A. Gostintsev, V.E. Fortov, Y.V. Shatskikh, Dokl. Phys. Chem., Part 1 **397**, 141–144 (2004)
46. L. Filyand, G.I. Sivashinsky, M.L. Frankel, On self-acceleration of outward propagating wrinkled flames. *Physica D* **72**, 110–118 (1994)
47. Y.C. Lee, H.H. Chen, Nonlinear dynamical models of plasma turbulence. *Phys. Scr. (Sweden)* **T2A**, 41–47 (1982)
48. O. Thual, U. Frisch, M. Henon, Application of pole decomposition to an equation governing the dynamics of wrinkled flame fronts. *J. Physique* **46**, 1485–1494 (1985)
49. G. Joulin, On the hydrodynamic stability of curved premixed flames. *J. Phys. Fr.* **50**, 1069–1082 (1989)
50. G. Joulin, On the Zhdanov-Trubnikov equation for premixed flame instability. *J. Exp. Theor. Phys.* **100**(2(8)), 428–432 (1990)
51. S.S. Minaev, Set of steady solutions describing a cellular flame in the case of hydrodynamic instability. *Combust. Explos. Shock Waves* **28**(1), 35–39 (1992)
52. S.S. Minaev, E.A. Pirogov, O.V. Sharypov, Nonlinear model of hydrodynamic instability of outward propagating flame. *Combust. Explos. Shock Waves* **32**(5), 481–488 (1996)
53. S.I. Glazyrin, Investigation of burning in type Ia supernovae. *Astron. Lett.* **39**(4), 221–226 (2013)
54. B. Shraiman, D. Bensimon, *Phys. Rev.* **A30**, 2840 (1984)
55. S.D. Howison, *J. Fluid Mech.* **167**, 439 (1986)
56. M. Rahibe, N. Aubry, G.I. Sivashinsky, R. Lima, Formation of wrinkles in outwardly propagating flames. *Phys. Rev. E* **52**(4), 3675–3686 (1995)
57. M. Rahibe, N. Aubry, G.I. Sivashinsky, Stability of pole solution for planar propagating flames. *Phys. Rev. E* **54**(5), 4958–4972 (1996)
58. M. Rahibe, N. Aubry, G.I. Sivashinsky, Instability of pole solutions for planar propagating flames in sufficiently large domains. *Combust. Theory Model.* **2**(1), 19–41 (1998)
59. Wm.T. Ashurst, Darrieus-Landau instability, growing cycloids and expanding flame acceleration. *Combust. Theory Model.* **1**, 405–428 (1997)
60. Y. Kortsarts, I. Brailovsky, G.I. Sivashinsky, On hydrodynamic instability of stretched flames. *Combust. Sci. Technol.* **123**, 207–225 (1997)
61. V. Karlin, G. Sivashinsky, The rate of expansion of spherical flames. *Combust. Theory Model.* **10**(4), 625–637 (2006)
62. V. Karlin, G. Sivashinsky, Asymptotic modelling of self-acceleration of spherical flames. *Proc. Combust. Inst.* **31**(1), 1023–1030 (2007)
63. R.V. Fursenko, K.L. Pan, S.S. Minaev, Noise influence on pole solutions of the Sivashinsky equation for planar and outward propagating flames. *Phys. Rev. E* **78**(5), 056301 (2008)
64. K.L. Pan, R.V. Fursenko, Characteristics of cylindrical flame acceleration in outward expansion. *Phys. Fluids* **20**(9), 094107 (2008)
65. R.V. Fursenko, S.S. Minaev, K.L. Pan, Hydrodynamic instability of inward-propagating flames. *Combust. Explos. Shock Waves* **45**(5), 511–517 (2009)
66. G. Joulin, P. Cambray, On a tentative, approximate evolution equation for markedly wrinkled premixed flames. *Combust. Sci. Technol.* **81**, 243 (1992)
67. P. Cambray, G. Joulin, Length-scales of wrinkling of weakly-forced, unstable premixed flames. *Combust. Sci. Technol.* **97**, 405–428 (1994)
68. P. Cambray, K. Joulain, G. Joulin, Mean evolution of wrinkle wavelengths in a model of weakly-turbulent premixed flame. *Combust. Sci. Technol.* **103**, 265 (1994)
69. G. Joulin, Nonlinear hydrodynamic instability of expanding flames: intrinsic dynamics. *Phys. Rev. E* **50**(3), 2030 (1994)
70. G. Borot, B. Denet, G. Joulin, Resolvent methods for steady premixed flame shapes governed by the Zhdanov-Trubnikov equation. *J. Stat. Mech.* **10**, 10023 (2012)
71. G. Joulin, B. Denet, Flame wrinkles from the Zhdanov-Trubnikov equation. *Phys. Lett. A Elsevier* **376**(22), 1797–1802 (2012)
72. G. Joulin, B. Denet, Shapes and speeds of steady forced premixed flames. *Phys. Rev. E Am. Phys. Soc.* **89**(6), 063001 (2014)

73. G. Joulin, B. Denet, Sivashinsky equation for corrugated flames in the large-wrinkle limit. *Phys. Rev. E* **78**, 016315 (2008)
74. B. Denet, Stationary solutions and Neumann boundary conditions in the Sivashinsky equation. *Phys. Rev. E* **74**, 036303 (2006)
75. L. Guidi, D. Marchetti, *Phys. Lett. A* **308**, 162 (2003)
76. M. Feigenbaum, I. Procaccia, B. Davidovich, *J. Stat. Phys.* **103**, 973–1007 (2001)
77. B. Denet, G. Joulin, Wrinkled flames and geometrical stretch. *Phys. Rev. E* **84**, 016315 (2011)
78. B. Denet, G. Joulin, Premixed-flame shapes and polynomials. *Physica D: Nonlinear Phenom.* vol. 292–293 (2015), pp. 46–50
79. L. Landau, *Acta Physicochimica U.R.S.S.* **19**, 77 (1944)
80. W.T. Ashurst, *Combust. Theory Model.* **1**, 405–428 (1997)
81. R. Aldredge, B. Zuo, *Combust. Flame* **127**, 2091–2101 (2001)
82. S.I. Blinnikov, P.V. Sasorov, *Phys. Rev. E* **53**, 4827–4841 (1996)
83. V.V. Bychkov, M.V. Liberman, *Phys. Rev. Lett.* **76**, 2814–2817 (1996)
84. A. Gostintsev, A.G. Istratov, V.E. Fortov, On fractal structure of turbulent spherical flame. *Dokl. RAN* **353**(1), 55–56 (1997)
85. K. Mukaiyama, S. Shibayama, K. Kuwana, Fractal structures of hydrodynamically unstable and diffusive-thermally unstable flames. *Combust. Flame* **160**, 2471–2475 (2013)
86. S. Gutman, G.I. Sivashinsky, The cellular nature of hydrodynamic instability. *Physica D* **43**, 129–139 (1990)
87. D. Bessis, J.D. Fournier, *J. Phys. Lett.* **45**, L-833 (1984)
88. F. Creta, N. Fogla, M. Matalon, Turbulent propagation of premixed flames in the presence of Darrieus-Landau instability. *Combust. Theory Model.* **15**(2), 267–298 (2011)
89. D. Vaynblat, M. Matalon, Stability of pole solutions for planar propagating flames: I. Exact eigenvalues and eigenfunctions. *SIAM J. Appl. Math.* **60**(2), 679–702 (2000)
90. D. Vaynblat, M. Matalon, Stability of pole solutions for planar propagating flames: II. Properties of eigenvalues/eigenfunctions and implications to stability. *SIAM J. Appl. Math.* **60**(2), 703–728 (2000)
91. B. Galanti, P.L. Sulem, A.D. Gilbert, *Physica D* **47**, 416 (1991) (and in the references therein)
92. H. Risken, *The Fokker-Planck Equation* (Springer, Berlin, 1984), p. 124 (Eqs. (5.110–5.111))
93. M.S. Uberoi, Flow field of a flame in a channel. *Phys. FL* **2**, 72 (1959)
94. B. Karlovitz, et al., A. Flame propagation across velocity gradients. B. Turbulence measurement in flames, in *Fourth Symposium (International) on Combustion*, Williams and Wilkins, Baltimore (1953), p. 613
95. A.G. Istratov, M. Gremyachkin, On the stability of plane flame in the flow with the velocity gradient. *Combustion and Explosion* (Nauka, Moscow, 1972), pp. 305–308 (In Russian)
96. G. Markstein, Experimental and theoretical studies of flame front stability. *Aero Sci.* **18**, 199 (1951)
97. G. Markstein, Instability phenomena in combustion waves. *Symp. (International) Combust.* **4**(1), 44 (1953)
98. H.F. Coward, F.J. Hartwell, Studies in the mechanism of flame movement. Part I. The uniform movement of flame in mixtures of methane and air, in relation to tube diameter. *J. Chem. Soc.* 1996–2004 (1932)
99. H.F. Coward, F.J. Hartwell, Studies in the mechanism of flame movement. Part II. The fundamental speed of flame in mixtures of methane and air. *J. Chem. Soc.* 2676–2684 (1932)
100. J. Quinard, G. Searby, B. Denet, J. Grana-Otero, Self-turbulent flame speeds. *Flow Turbul. Combust.* **89**, 231–247 (2012)
101. M. Kuznetsov, M. Liberman, I. Matsukov, Experimental study of the preheat zone formation and deflagration to detonation transition. *Combust. Sci. Technol.* **182**, 1628–1644 (2010)
102. L. Kagan, G. Sivashinsky, The transition from deflagration to detonation in thin channels. *Combust. Flame* **134**(4), 389–397 (2003)
103. V. Karlin, Radiation preheating can trigger transition from deflagration to detonation. *Flow Turbul. Combust.* **87**(2–3), 511–523 (2011)

104. O. Kupervasser, Pole solutions in the case of problems of flame front propagation and Saffman-Taylor “finger” formation without surface tension: open problems and possible ways of their solutions (2003), <http://arxiv.org/abs/nlin/0306038>
105. N. Fogla, Interplay between background turbulence and Darrieus-Landau instability in premixed flames via a model equation. Ph.D. thesis, University of Illinois at Urbana-Champaign, Urbana, Illinois (2010)
106. T.A. Witten, L.M. Sander, *Phys. Rev. Lett.* **47**, 1400 (1981)
107. P. Meakin, *Phys. Rep.* **235**, 189 (1993)
108. T. Halpin-Healey, Y.C. Zhang, *Phys. Rep.* **254**, 215 (1995)
109. Y. Kuramoto, *Suppl. Prog. Theor. Phys.* **64**, 346 (1978)
110. V.S. L'vov, I. Procaccia, *Phys. Rev. Lett.* **69**, 3543 (1992)
111. M.-H. Wu, W.-C. Kuo, Accelerative expansion and DDT of stoichiometric ethylene/oxygen flame rings in micro-gaps. *Proc. Combust. Inst.* **34**(2), 2017–2024 (2013)
112. D.A. Kessler, J. Koplik, H. Levine, Pattern selection in fingered growth phenomena. *Adv. Phys.* **37**, 255–339 (1986)
113. D. Khavinson, M. Mineev-Weinstein, M. Putinar, Planar elliptic growth. *Complex Anal. Oper. Theory* **3**, 425–451 (2009)
114. D. Khavinson, M. Mineev-Weinstein, M. Putinar, R. Teodorescu, Lemniscates do not survive Laplacian growth. *Math. Res. Lett.* **17**(2), 337–343 (2010)
115. E. Lundberg, V. Totik, Lemniscate growth. *Anal. Math. Phys.* **3**, 45–62 (2013)
116. E. Lundberg, Laplacian growth, elliptic growth, and singularities of the Schwarz potential. *J. Phys. A: Math. Theor.* **44**, 135202 (2011)
117. N.N. Smirnov, V.F. Nikitin, A. Maximenko, M. Thiercelin, J.C. Legros, Instability and mixing flux in frontal displacement of viscous fluids from porous media. *Phys. Fluids* **17**, 084102 (2005)
118. V.R. Dushin, V.F. Nikitin, Yu.G. Philippov, N.N. Smirnov, Two phase flows in porous media under microgravity conditions. *Microgravity Sci. Technol.* **20**(3–4), 155–160 (2008)
119. N.N. Smirnov, V.F. Nikitin, V.R. Dushin, A. Maximenko, M. Thiercelin, J.C. Legros, Instability in displacement of viscous fluids from porous specimens. *Acta Astronautica* **61**(7–8), 637–643 (2007)
120. Ar. Abanov, M. Mineev-Weinstein, A. Zabrodin, Multi-cut solutions of Laplacian growth. *Physica D: Nonlinear Phenom.* **238**(17), 1787–1796 (2009)
121. S.Y. Lee, R. Teodorescu, P. Wiegmann, Shocks and finite-time singularities in Hele-Shaw flow. *Physica D: Nonlinear Phenom.* **238**(14), 1113–1128 (2009)
122. S.Y. Lee, R. Teodorescu, P. Wiegmann, Weak solution for the Hele-Shaw problem: viscous shocks and singularities. *JETP Lett.* **92**(2), 91–96 (2010)
123. S.J. Chapman, *Eur. J. Appl. Math.* **10**, 513–534 (1999)
124. R. Combescot, T. Dombre, V. Hakim, Y. Pomeau, *Phys. Rev. Lett.* **56**, 2036–2039 (1986)
125. R. Combescot, V. Hakim, T. Dombre, Y. Pomeau, A. Pumir, *Phys. Rev. A* **37**, 1270–1283 (1988)
126. D.C. Hong, J.S. Langer, *Phys. Rev. Lett.* **56**, 2032–2035 (1986)
127. B.I. Shraiman, *Phys. Rev. Lett.* **56**, 2028–2031 (1986)
128. S.J. Chapman, J.R. King, *J. Eng. Math.* **46**, 1–32 (2003)
129. S. Tanveer, *J. Fluid Mech.* **409**, 273–308 (2000)
130. L.A. Galin, Unsteady filtration with a free surface. *Dokl. Akad. Nauk USSR* **47**, 246–249 (1945) (in Russian)
131. P. Ya, Polubarinova-Kochina, On a problem of the motion of the contour of a petroleum shell. *Dokl. Akad. Nauk USSR* **47**, 254–257 (1945) (in Russian)
132. P.P. Kufarev, A solution of the boundary problem for an oil well in a circle. *Dokl. Akad. Nauk USSR (N.S.)* **60**, 1333–1334 (1948) (in Russian)
133. M. Siegel, S. Tanveer, *Phys. Rev. Lett.* **76**, 409–422 (1996)
134. A. Sarkissian, H. Levine, *Phys. Rev. Lett.* **81**, 4528 (1998)
135. M. Mineev-Weinstein, A reply to the comment by Armand Sarkissian and Herbert Levine. *Phys. Rev. Lett.* **81**, 4529–4529 (1998)

136. J. Casademunt, F.X. Magdaleno, Comment on “Selection of the Saffman-Taylor finger width in the absence of surface tension: an exact result”. *Phys. Rev. Lett.* **81**, 5950–5950 (1998)
137. R.F. Almgren, Comment on “Selection of the Saffman-Taylor finger width in the absence of surface tension: an exact result”. *Phys. Rev. Lett.* **81**, 5951–5951 (1998)
138. M. Mineev-Weinstein, A reply to the comment by J. Casademunt and F. X. Magdaleno, and also R.F. Almgren. *Phys. Rev. Lett.* **81**, 5952–5952 (1998)
139. D.A. Kessler, H. Levine, Microscopic selection of fluid fingering patterns. *Phys. Rev. Lett.* **86**, 4532–4535 (2001)
140. E. Paune, F.X. Magdaleno, J. Casademunt, Dynamical systems approach to Saffman-Taylor fingering. A dynamical solvability scenario. *Phys. Rev. E* **65**(5), 056213 (2002)
141. M. Mineev-Weinstein, G.D. Doolen, J.E. Pearson, S.P. Dawson, Formation and pinch-off of viscous droplets in the absence of surface tension: an exact result, <http://arxiv.org/abs/patt-sol/9912006>
142. A.P. Aldushin, T.P. Ivleva, Modelirovanie gidrodinamicheskoy neustojchivosti filtracionnogo rezhima rasprostraneniya fronta gorenija v poristoj srede. Ph.D. thesis, Institut strukturnoj makrokinetiki i problem materialovedeniya RAN, Chernogolovka, Russia (in Russian)
143. A.P. Aldushin, B.Sh. Braverman, Hydrodynamic instability of filtration combustion wave. *Dokl. Phys. Chem., Part 1* **427**, 340–343 (2009)
144. O. Zik, Z. Olami, E. Moses, Fingering instability in combustion. *Phys. Rev. Lett.* **81**, 3868 (1998)
145. O. Zik, E. Moses, Fingering instability in combustion: an extended view. *Phys. Rev. E* **60**(1), 518 (1999)
146. A.P. Aldushin, B.S. Seplyarskii, *Dokl. Akad. Nauk SSSR* **241**, 72 (1978) (*Sov. Phys. Dokl.* **23**, 483 (1978))
147. A.P. Aldushin, *Prog. Aeronaut. Astronaut. AIAA* **173**, 95 (1997)
148. P.G. Saffman, G.I. Taylor, *Proc. R. Soc. A* **245**, 312 (1958)
149. G. Taylor, P.G. Saffman, *Q. J. Mech. Appl. Math.* **12**, 265 (1959)
150. J.M. Vanden-Broeck, *Phys. Fluids* **26**, 2033 (1983)
151. A.P. Aldushin, T.P. Ivleva, Continuous infiltration mediated SHS process in a coflow reactor: numerical modeling. *Int. J. Self Propag. High Temp. Synth.* **22**(2), 77–83 (2013)
152. A.P. Aldushin, T.P. Ivleva, Hydrodynamic instability of the coflow filtration combustion: numerical simulation. *Dokl. Phys. Chem., Part 1* **451**, 157160 (2013)
153. A.P. Aldushin, T.P. Ivleva, Coflow filtration combustion in a slit-like reactor: simultaneous manifestation of fluid dynamic and thermal instability. *Int. J. Self Propag. High Temp. Synth.* **23**(3), 121–532 (2014)

Index

C

Cusp, 2–4, 9, 13, 14, 19, 21, 24, 25, 29–31, 36, 37, 42, 45, 46, 68–78

D

Darrieus-Landau hydrodynamic instability, 4, 5, 45, 77, 78
Diffusional-thermal instability, 80

F

Filtration combustion, 2, 85–87, 102, 103, 105–107
Finite time singularity, 86–88, 90, 91, 97, 98, 100, 101, 104
Front propagation, 1, 3, 6, 9–11, 47, 67, 78, 81, 98, 102, 104, 109

G

Giant cusp, 2, 4, 6–9, 12, 17–21, 25, 27, 30–32, 35, 36, 39–43, 46, 58–65, 71–74, 76

H

Hele-Shaw cell, 2, 86, 96
Hele-Shaw channel, 85
Hydrodynamic instability, 1, 2, 4, 80, 103

I

Interface growth, 1, 68

L

Laplacian growth, 1–4, 6, 9, 10, 47, 68, 71, 72, 83, 85–90, 97, 98, 102, 104, 105, 109, 110

M

Michelson-Sivashinsky equation, 4

N

Navier–Stokes equations, 1

P

Pole solution, 2, 4, 7, 9, 10, 18, 35, 40, 41, 85, 90, 98, 101, 104
Premixed flame, 1–6, 8–13, 20, 23–25, 27–30, 43–52, 55, 65, 67, 68, 71, 72, 74, 75, 77–83, 86, 87, 98, 104, 109

Q

Quasi-cusp, 4, 14, 45, 70, 78

S

Saffman-Taylor “finger”, 88, 97, 98, 101
Saffman-Taylor (SF) problem, 107
Saffman-Taylor (ST) problem, 87
Self-acceleration, 1–4, 77, 80, 81
Sivashinsky, Filyand and Frankel equation, 2, 81

Stability analysis, [9](#), [10](#), [12](#), [15](#), [16](#), [18](#), [35](#),
[55–58](#), [65](#), [73](#), [109](#)
Surface tension, [2](#), [3](#), [6](#), [9](#), [10](#), [86–88](#), [97](#), [98](#),
[101](#), [102](#), [104](#), [107](#), [110](#)

T

TFH-cusp, [8](#), [15](#), [71](#), [73](#)
TFH-giant cusp, [73](#)
TFH solution, [61](#)

Mathematical and Analytical Techniques with Applications to Engineering

Oleg Kupervasser

Pole Solutions for Flame Front Propagation

This book deals with solving mathematically the unsteady flame propagation equations. New original mathematical methods for solving complex non-linear equations and investigating their properties are presented. Pole solutions for flame front propagation are developed. Premixed flames and filtration combustion have remarkable properties: the complex nonlinear integro-differential equations for these problems have exact analytical solutions described by the motion of poles in a complex plane. Instead of complex equations, a finite set of ordinary differential equations is applied. These solutions help to investigate analytically and numerically properties of the flame front propagation equations.

Engineering

ISBN 978-3-319-18844-7



► springer.com